AFRICAN SWINE FEVER (ASF) Situation Report 59

Period covered: November 2024

This report provides an update of the ASF situation, according to the information shared with WOAH.

Key highlights

- During the period covered by this report, **one country in Europe and one territory in Asia** reported new ASF events, while **nine countries in Europe** updated their ongoing events. No new outbreak was reported by countries/territories in Africa, the Americas and Oceania. **Twenty-five new outbreaks** were reported in domestic pigs and **197** in wild boar, in Asia and Europe, with **602 animal losses in domestic pigs**.
- Most of the outbreaks reported during the period are in high density pig farming areas.
- The number of outbreaks reported in domestic pigs and wildlife through immediate notifications and follow up reports has been globally declining since July 2024, but 10 outbreaks were reported more than 10 km outside previously affected areas, up to 75 km in Albania.
- Since January 2022, **11 countries** have reported ASF as a first occurrence in the country, while **11 countries** have reported its spread to new zones.
- Since January 2022, more than **741,000 cases** in pigs and more than **26,400 cases** in wild boars have been reported, with more than **1,795,000 animal losses in domestic pigs**.
- Since January 2022, **63 countries and territories (62 countries and territories until last update)** have reported the presence of ASF.



Contextual information of the ASF situation by world region (1 January 2022 – 31 October 2024)

In total, during the period, ASF has been reported as present in five different world regions in 63 countries, affecting more than 741,000 pigs and more than 26,400 wild boars, with more than 1,794,000 animal losses. Further details, split by world region, are included in Table 1. During the period, no country/territory reported vaccination of pigs in response to the outbreaks.

Table 1. Summary of the number of outbreaks, cases and animal losses caused by ASF in the different world regions since January 2022.

	Outbreaks		Cases		Losses*	
	Domestic pigs	Wild boar	Domestic pigs	Wild boar	Domestic pigs	
Africa	282	5	79,635		68,696	
Americas	53	0	355	0	4,940	
Asia	4,089	106	219,504	539	427,204	
Europe	4,337	15,781	442,163	25,885	1,294,095	
Oceania	0	0	0	0	0	
Total	8,761	15,892	741,657	26,424	1,794,935	

*Losses (deaths + animals killed and disposed of): this figure refers to losses in the establishments affected by the outbreaks and it does not include the animals culled in areas around the outbreak for controlling the disease.

The spatial distribution of outbreaks reported since January 2022 in domestic pigs and wildlife is shown in Figure 1.




Figure 1. Map of ASF outbreaks which started during 01 Jan 2022 – 31 Oct 2024 in domestic pigs and wildlife.

Recent updates (01/11/2024 – 30/11/2024)

To describe the current disease situation of ASF, this section covers: (a) a list of new events which started during the period (reported through INs); (b) information on events that started before the period but were still ongoing during the period (reported through FURs); and (c) the geographic distribution of new outbreaks that started during the period. This information is based on immediate notifications (INs) and follow up reports (FURs) received by the World Organisation for Animal Health (WOAH) through the World Animal Health Information System (WAHIS). The outbreaks are displayed on a map in Figure 1.

New events by world region (reported through INs, see Figure 3)

<u>Asia</u>

Hong Kong (SAR-PRC) reported the recurrence of the disease (the event started on 21 November)

<u>Europe</u>

Albania reported the recurrence of the disease (the event started on 25 November in Lezhë)

Africa, Americas, and Oceania

No new events reported

On-going events for which there were new outbreaks, by world region (reported through FURs, see Figure 3)

<u>Europe</u>

Bosnia and Herzegovina, Germany, Hungary, Italy, North Macedonia, Poland, Romania, Serbia, and Ukraine.

Africa, Americas, Asia, and Oceania

No ongoing events updated

The number of outbreaks, cases and losses during the period covered by this report are displayed in Table 2. During the period covered by the report, no country/territory reported vaccination of pigs in response to the outbreaks.

Table 2. Summary of the number of outbreaks, cases and animal losses caused by ASF in the different world regions during the reporting period.

	Outbreaks		Cases		Losses*
	Domestic pigs	Wild boar	Domestic pigs	Wild boar	Domestic pigs
Africa	0	0	0	0	0
Americas	0	0	0	0	0
Asia	0	1	0	1	0
Europe	25	196	1,463	250	602
Oceania	0	0	0	0	0
Total	25	197	1,463	251	602

*Losses (deaths + animals killed and disposed of): this figure refers to losses in the establishments affected by the outbreaks and it does not include the animals culled in areas around the outbreak for controlling the disease.

The distribution of outbreaks is shown in Figure 3. To highlight the impact of disease spread on the pig industry, the density of pigs is shown in the background. Most of the outbreaks reported during the period are in high-density pig farming areas.

If we take the geographical distribution of the disease between 1 January and 31 October 2024 as a reference, we note that during the period covered by this report, 10 outbreaks were notified more than 10 km outside the previous geographical distribution. The most distant outbreak (in Albania) was reported 75 km from the previous geographical distribution.

Regarding the temporal dynamics of the disease as reported through the WAHIS early warning system (excluding areas with stable situations), Figure 2 shows the evolution of the monthly number of reported outbreaks in domestic and wild animals since 2022 and as of 30 November (taking into account both INs and FURs). Both the domestic and wildlife trends show an overall decreasing trend in the number of reported outbreaks.




Figure 2. Trends in the monthly number of reported ASF outbreaks in domestic and wild animals for the period 01 Jan 2022 – 30 Nov 2024 reported through the WAHIS early warning system (excluding endemic areas).




Figure 3. Map of ASF outbreaks which started between 01 Nov 2024 and 30 Nov 2024 in domestic animals and wildlife. Zoomed views are provided as well. The density of pigs based on <u>FAO GLW 4: Gridded</u> <u>Livestock Density</u> (as of 2020) is shown in the background in shades of blue.

Self-declaration of freedom from ASF submitted during the reporting period

Eighteen self-declarations from 17 countries are currently active and can be consulted in the <u>dedicated dashboard</u> on WOAH website.

During the reporting period, no new self-declaration of freedom from ASF was publishedby WOAH.

Recommendations

- ASF continues to represent a global threat, and WOAH highlights the importance of implementing strict biosecurity, an early reporting and response system, while maintaining a high level of disease awareness among all actors involved in the value chain.
- There are countries that have approved or are conducting field trials of the use of modified live vaccine candidates against ASF Genotype II. As with all vaccines, <u>WOAH stresses the importance of using only high-quality vaccines</u> with demonstrated effectiveness and safety, in accordance with standards in the *Terrestrial Manual*, including those that have been drafted for ASF vaccines.
- As of 30 November 2024, no countries and territories have officially reported to WOAH the implementation of vaccination in response to ASF around reported outbreaks. WOAH urges Members who have a vaccination programme in place to share the information with WOAH and the international community.
- Any vaccination strategy for ASF should be undertaken as part of a well-designed vaccination programme that considers factors including the local epidemiology of ASF, the circulating strains, the expected objectives and the adequacy and sustainability of the relevant technical, financial and human resources. The vaccination programme should also include post-vaccination surveillance and monitoring as well as an exit strategy for the cessation of vaccination, as per <u>Chapter 4.18</u>, of the *Terrestrial Code*.
- WOAH urges its Members to continue to promptly notify the occurrence of ASF and to share the relevant epidemiological information, including information on any newly detected recombinant strains and vaccination trials that can facilitate transparency and assist the global control of the disease.

More information and WOAH resources

- WOAH ASF webpage
- WOAH regional webpages for ASF which provides regional updates on the disease situation and activities:
 - <u>Africa</u>
 - <u>Americas</u>
 - Asia and the Pacific
 - Europe
- WOAH and FAO designed communication tools on ASF for use by any interested party
- WOAH Terrestrial Animal Health Code
- WOAH Manual of Diagnostic Tests and Vaccines for Terrestrial Animals
- ASF Reference Laboratory <u>summary</u> of available PoC kits to guide field workers, practitioners and decision-makers in their use and <u>laboratory algorithm manual</u> to address the detection of virulent and variant forms of ASFV.
- <u>Global Framework for the Progressive Control of Transboundary Animal Diseases (GF-TADs)</u> page for ASF
- Global African Swine Fever Research Alliance

For any press inquiry on ASF, e-mail us at media@woah.org.